This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

INTERNATIONAL APPLICATION FOREIG			
(51) International Patent Classification 4:		(11	1) International Publication Number: WO 88/ 06630
C12P 21/00, C07H 17/00 C07K 15/04, A61K 39/395	A1	(43	3) International Publication Date: 7 September 1988 (07.09.88)
(21) International Application Number: PCT/US (22) International Filing Date: 2 March 1988		- 1	(74) Agents: FOX, Samuel, L. et al.; Saidman, Sterne, Kessler & Goldstein, 1225 Connecticut Avenue, N.W., Suite 300, Washington, DC 20036 (US).
(31) Priority Application Number: (32) Priority Date: 2 March 1987 (33) Priority Country:			(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent).
(71) Applicant: GENEX CORPORATION [US/U Industrial Drive, Gaithersburg, MD 20877 (172) Inventors: LADNER, Robert, Charles; 382 Valley Road, Ijamsville, MD 27754 (US). G Leslie; 10899 Deborah Drive, Potomac, M (US). BIRD, Robert, E.; 3903 Morrell Co sington, MD 20895 (US).	US). 27 Gre LICK, 1D 208	en J., 54	With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt

(54) Title: METHOD FOR THE PREPARATION OF BINDING MOLECULES

(57) Abstract

This invention comprises a genetically engineered organism displaying the expression product of an inserted gene on its outer surface. In a preferred embodiment, a single chain antibody is displayed on the outer surface of the genetically engineered microorganism.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

					•
AT	Austria	FR	France	ML	Mali
ΑŪ	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JР	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM.	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark	MC	Monaco	US	United States of America
FI .	Finland	MG	Madagascar	-	

-1-

Method for the Preparation of Binding Kolecules

Background of the Invention

Field of the Invention

The present invention relates to the production of genetically engineered organisms and methods of preparation of binding molecules.

Description of the Background Art

In vertebrates, antibody diversity arises from substitution of hypervariable loops into constant antibody frameworks. Each B-cell exhibits its own type of specificity on its surface. When an antigen binds to the surface antibody, the B-cell is stimulated to proliferate.

Monoclonal antibody production exploits this as follows: An animal is injected with a purified antigen. After several weeks, the spleen is removed from the animal and spleen cells are fused to myeloma cells. This produces hybridoma cells. These cells are plated and screened for binding to antigen. These cells can be grown in tissue culture and will produce quantities of a single antibody—a monoclonal antibody.

The gene for the antibody can be recovered and put into microorganisms. Genetic and protein engineering can be altered to obtain better binding, altered specificity,

different antigenic behavior than that of the original protein or gene product.

Single-chain antibodies (SCA) (copending U.S. Patent, Application Serial Nos. 902,971 and 902,970, herein incorporated by reference) are protein molecules which retain the binding domain of antibodies but not the effector domains.

SUMMARY OF THE INVENTION

In the present invention, a genetically engineered organism is produced which displays on the outer surface of the organism the expression product of a gene which has been inserted.

In one embodiment of the invention, a SCA domain (SCAD) is displayed on the <u>outside</u> of a microorganism while the message for that particular SCA is <u>inside</u> that organism.

Description of the Drawings

Figure 1, flow chart for production of organisms containing binding molecules on surface.

Figure 2, displaying SCAD on surface.

Figure 3, making diverse population of displayed SCADs.

Figure 4, selecting new SCA specificity.

Figure 5, detecting known antigens.

Figure 6, lambda assembly.

Figure 7, inserting SCAD into V genes.

DESCRIPTION OF PREFERRED EMBODIMENTS

Any protein or antibody domain for which a gene can be isolated or constructed may be displayed on the outer surface of an organism into which the gene has been inserted. This is done by fusing the SCAD gene to the gene coding for a product which normally expresses on the surface of the organism; e.g., an envelope protein. The organism so

produced may be easily isolated from organisms which do not contain the desired gene and express the gene product. The organisms may also serve as a solid substrate for the gene product. Prior to the present invention, once an organism which contained a desired gene had been produced, the organisms had to be grown and assayed for the production of the gene product. Then, the gene product had to be isolated, purified, and only then was it possible to couple it to a solid substrate. In one embodiment, the organism itself containing the gene product on its outer surface is the solid substrate with the desired gene product already attached and may be used as such.

The present invention is depicted in general terms in the flow chart on Figure 1. One embodiment shown at step 1000 consists of producing organisms. In this embodiment, a microorganism displays a gene product such as a SCAD on the surface of the organism. The next step (step 1010) consists of generating, from the one SCAD displayed and encoded in the organism, a diverse population of SCADs by varying the DNA sequence encoding the SCAD by mutation techniques. The new diverse SCADs generated in step 1010 are displayed on the surface of the organism (step 1020) and organisms are selected based on the surface expressed SCAD which bind to given antigens (step 1030). The organisms selected in step 1030 may be used in assays for the given antigen or may be further selected according to the binding or enzymatic characteristics of the gene product expressed on the surface.

Once any SCAD has been displayed on the surface of a microorganism, a large population of <u>different</u> SCADs can be generated by <u>in vivo</u> DNA synthesis, step 1010, and each cell or virion can display its own SCAD specificity, step 1020.

Antigen binding to the displayed SCAD can be used to select

those microorganisms harboring genes for SCADs which will bind antigen, step 1030. Once a strain of microorganisms is selected for antigen binding, it can be used as a sensitive assay for that antigen, step 1040. In step 1050, the ability to refine antigen binding is used to generate novel enzymes.

The steps needed to achieve the construction of a microorganism which displays SCAD are shown in the flow chart of
Figure 2. In step 2000, a microorganism is selected. In
step 2010, a gene within that organism is selected; the gene
must be one coding for a protein which is displayed on the
cell or virion surface. Preferably, the gene should not be
essential to the organism. In step 2020, the gene for a SCAD
to some known antigen is introduced into the selected gene,
and in step 2030, this population of modified genes is put
back into the organism. In step 2040 the genes are expressed, and in step 2050 the organisms are selected for binding
to immobilized antigen. In step 2060, the gene is sequenced
to determine which insertion was fruitful.

The steps needed to create a diverse population of SCADs displayed on the surface of a microorganism are illustrated in the flow chart on Figure 3. In step 3000, the Combining Determining Regions (CDRs) of the SCA are bounded by restriction sites. In step 3010, a large variety of DNA sequences are produced. Each sequence should begin with one of the restriction sequences and end with the corresponding restriction site. Between these sites should come any constant residues which are included to facilitate placement of restriction sites plus an integral number of triplets. The number of triplets can be varied within the bounds set by:

- 1. Analysis of sequences of natural antibodies with similar framework;
 - Computer modeling of the framework;
 - Trial and error.

In step 3020, these DNA sequences are inserted into the appropriate slots in the SCAD gene. In step 3030, these genes are reinserted into the organism and grown. The organism now contains a diversity of SCA specifications, each cell displaying its own particular SCAD. In step 3040, this population is passed over the inert support which will be used to support antigen. This step removes those organisms which bind to the inert support even without antigen.

In step 4000 in Figure 4, the antigen is attached to an inert support. In step 4010, the population of organisms prepared in steps 3000 to 3040 is passed over the supported antigen. Organisms not binding pass through. In step 4020, the organisms bound to the support are allowed to grow. In step 4030, colonies are found and sampled. In step 4040, the genes of several isolates are sequenced. In step 4050;, the SCAD gene of selected organisms are mutagenized. In step 4060, step 4010 through 4050 are repeated with the mutagenized colonies. In step 4060, by washing more stringently, a SCA colony with maximal binding is obtained. Step 4060 can be repeated until suitable binding is obtained.

The present invention is also useful for the detection and quantification of known antigens. In step 5000 of Figure 5, a sample with unknown amount of an antigen is attached to an inert support. In step 5010, the strain of organism derived in step 4060 and displaying a SCAD against the antigen is passed over the inert support. In step 5020, the bound organisms are allowed to grow. In step 5030, points of growth are detected, the amount of growth quantitates the amount of antigen.

Enzymes, particularly degradative enzymes, work by stabilizing the transition state of a reaction. Chemical theory suggests the shape of the transition states of many reactions. For example, the carbonyl carbons of esters of

carboxylic acids are trigonal planar. The transition state for hydrolysis or transesterification is almost certainly tetrahedral. It has recently been demonstrated that a monoclonal antibody against a phosphate ester (which is tetrahedral) is also an esterase.

Monoclonal antibody technology has many shortcomings for this task:

- 1. Slow turnaround
- 2. Difficulty in refining antibody
- 3. Inappropriate for highly toxic chemicals
- 4. Inappropriate for metabolites

The method described above can be applied to produce such enzyme-like binding molecules. It will be limited only in the ability to invent antigens which resemble the transition states of reactions wished to catalyze.

For example, to catalyze the transesterification

one can raise a SCA against

Having now generally described the invention, the same may further be understood by reference to the following

expressly stated.

Example 1

The preferred embodiment utilizes the bacteriophage lambda. The gene V of lambda generates a protein which assembles to form the neck of lambda. First gene product V (gpV) forms hexameric annuli, then 32 of these annuli stack on the nose cone to form the neck (Figure 6). Finally, the neck joins the head which contains the DNA.

gpV is a protein of molecular weight 31K. Wild-type lambda have small protuberances on the outside of the neck annuli. Mutants have been isolated in which as much as 13K of gpV is absent. These mutants are viable, though temperature sensitive. The mutants are those wherein shortened gpV lack the protuberances on the neck annuli. Genetics indicates that the deletion is from the carboxy end of gene V.

SCAs made so far contain the four cysteine residues found in all VH and Vl domains of natural antibodies.

Natural antibodies are secreted and fold in the oxidizing environment of serum. The interior of cells is a reducing environment; thus, one would not expect disulfide bonds to form. The sulfhydryl groups of cysteine lie only 2.0 A apart when a disulfide forms. If the disulfide is reduced, the sulfur atoms should lie 4.0 A apart. Thus, reduced cysteines will greatly destabilize folding of a SCA. Therefore, to get proper folding of SCAD inside a cell, one mutates the SCAD gene to change all or some of the CYS's to SER, THR, ALA, or GLY. In one embodiment, the SCA is against bovine growth hormone (BGH).

The V gene of lambda is shown in Figure 7. Genetics indicates that the domain responsible for the warts on the neck lies in the 300 to 400 last base pairs to the right. One cuts the gene at some point in this region, preferably

200 bases from the right end. A random number of bases on either side, up to 200 bases is removed. The SCAD (antiBGH) is inserted and put back into a lysogenic strain of <u>E. coli</u>. In the preferred embodiment, the lambda contains a highly beneficial gene for the <u>E. coli</u>.

The <u>E. coli</u> is induced. The lambda progeny is passed over a support holding BGH. The <u>E. coli</u> is contacted with the support. The <u>coli</u> should be deficient in a way that the beneficial gene in the lambda will complement. For example the <u>coli</u> could be drug-sensitive and lambda will carry drug resistance. The corresponding antibiotic in the medium puts the <u>coli</u> under selective pressure so that only those cells infected by lambda will grow. Only those lambda which bound antigen and stuck to the support are available.

WE CLAIM:

- 1. An organism containing a recombinant gene wherein the product of said recombinant gene is presented on the outer surface of said organism, said product also comprising a single chain antibody domain (SCAD).
- 2. The organism of claim 1 wherein the product of said recombinant gene comprises a polypeptide which directs said SCAD to the surface of said organism.
- 3. A fusion polypeptide comprising a product normally appearing on the surface of an organism fused to a single chain antibody domain.
- 4. A method of preparing an organism containing a single chain binding molecule on the outer surface of said organism which comprises:
- (1) isolating from an organism a first gene encoding for a cell surface protein;
- (2) inserting a second gene which encodes a single chain antibody domain into said first gene to form a recombinant fusion gene; and
- (3) transforming an organism with said recombinant fusion gene.




FIG. 1

2/7

2000		
Select a microorganism		
2010		
Select gene for surface protein		
2020		
Select SCAD (Anti Known Antigen)		
2030		
Insert SCAD gene into surface gene ;		
2040		
Return surface gene to organism		
2050		
Select for binding antigen		
2060		
Sequence selected gene		

FIG. 2

3000			
	Delimit CDRs of SCAD by		
restriction of s	restriction of sites		
3010			
Produce, in vitro DNA sequences	, large variety of for CDR's		
3020			
Insert DNA pieces into CDR slots			
3030			
Reinsert genes i	nto organism		
3040			
Deplete populati			
to inert support			

FIG. 3




FIG. 4




FIG. 5 Detecting Known Antigens






FIG. 6

SUBSTITUTE SHEET








FIG. 7

INTERNATIONAL SEARCH REPORT

International Application NPCT/US 88/00716

I. CLASSIFICATION OF SUBJECT MATTER (II several classification sympols apply, indicate all) 3						
According to International Patent Classification (IPC) or to both National Classification and IPC						
1						
C12P 21/00, C07H 17/00, C07K 15/04, A61K 39/395						
II. FIELDS SEARCHED						
	Minimum Documentation Searched 4					
Classification Syst	em Classification Symbols					
US	435/68, 70, 172.3, 235, 948, 91, 530/387	, 388,				
	424/85, 536/27					
	Documentation Searched other than Minimum Documentation					
	to the Extent that such Documents are included in the Fields Searched 6					
Computer	Search CAS, BIOSIS, APS: Immunogloulin,	clone 1,				
plasmid	1, surface, secrete, signal or leader, coa-	t protein,				
	ctor, GPV protein					
	S CONSIDERED TO BE RELEVANT 14					
	Citation of Document, 16 with indication, where appropriate, of the relevant passages 17	Relevant to Claim No. 18				
	ence, Vol. 81, issued August, 1987,	1-4				
(Wa	shington, D.C., U.S.A.), (R. HUBER)					
	ructural Basis for antigen-antibody					
	ognition". See entire document.					
	es 702-703.					
Y U.S	U.S.,A, 4,704,692, (LADNER) issued 03					
Nov	ember 1987. See the entire document.					
Y U.S	.,A, 4,603, 112, (PAOLETTI ET AL)	1-4				
iss	ued 29 July 1986. See columns 1-3					
in	particular and abstract.					
Y Bio	technology, Vol. 3, issued April	1-4				
198	5, (New York, New York, U.S.A.),	ļ				
(VA	LENZUELA ET AL) "Antigen Engineering	l				
in	Yeast: Synthesis and Assembly of					
Hyb	rid Hepatitis B Surface Antigen-Herpes	•				
Sim	plex 1 gD Particles", pages 323-326.					
1						
I						
• Special estate	ros of cital documents: 15 "T" later document nublished after the	international filles date				
*Special categories of cited documents: 15 "A" document defining the general state of the art which is not or priority date and not in conflict with the application but or priority date and not in conflict with the application but or priority date and not in conflict with the application but or priority date and not in conflict with the application but						
considered (to be of particular relevance invention					
"E" earlier document but published on or after the international filing date "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to						
"L" document which may throw doubts an priority claim(s) or involve an inventive step						
citation or other special reason (as specified) Cannot be considered to involve an inventive step when the						
"O" document referring to an oral disclosure, use, exhibition or document is combined with one or more other such document other means. Such combination being obvious to a person skilled						
"P" document published prior to the international filing date but						
later than the priority date claimed "&" document member of the same patent family						
IV. CERTIFICATION						
Date of the Actual Completion of the International Search Date of Mailing of this International Search Report						
2 4 JUN 1988						
06 MAY 1988 International Searching Authority 1 Signature of Authorized Officer 10						
	DJ. H. Huba					
ISA/US	ROBIN L. TESKIN					

Form PCT/ISA/210 (second sheet) (October 1981)

Category •	Citation of Document, In with indication, where appropriate, of the relevant passages of	Relevant to Claim No I
A	Biotechnology, Volume 4, issued April 1986, (New York, New York), (J. Van Brunt) "Protein Architecture: Design-ing from the Ground Up", see pages 277-283.	1-4
Y .	U.S.A., 4,593,002, (DULBECCO) issued 03 June 1986. See the entire document.	1-4
·		
·		
	·	